

# Enzymology: General Concepts and Enzyme Kinetics

---

## Enzymology: General Concepts and Enzyme Kinetics

---

### Topics: Classification, Co-enzymes

(No figures, clean text-only style.)

---

### Classification of Enzymes

---

Enzymes are classified by the **International Union of Biochemistry (IUB)** into **six major classes** based on the reactions they catalyze.

#### 1. Oxidoreductases

- Catalyze **oxidation–reduction** reactions.
  - Transfer of electrons or hydrogen atoms.
  - Examples: **Dehydrogenases, oxidases, reductases.**
- 

#### 2. Transferases

- Transfer **functional groups** (methyl, amino, phosphate).
  - Examples: **Transaminases, kinases, methyltransferases.**
- 

#### 3. Hydrolases

- Catalyze **hydrolysis** of bonds (using water).
  - Examples: **Proteases, lipases, amylases, phosphatases.**
- 

#### 4. Lyases

- Break bonds **without hydrolysis or oxidation**, forming double bonds.
  - Examples: **Decarboxylases, aldolases.**
-

## 5. Isomerases

- Catalyze **intramolecular rearrangements**.
  - Examples: **Racemases, epimerases, mutases**.
- 

## 6. Ligases (Synthetases)

- Join two molecules together using **ATP**.
  - Examples: **Carboxylase, DNA ligase**.
- 

### Mnemonic

**O-T-H-L-I-L** ? *“Only Tigers Hunt Lions In Laos”*

---

## Additional Enzyme Classification Concepts

---

### Based on Composition

- **Simple enzymes** ? made of protein only.
  - **Conjugated enzymes** ? protein (apoenzyme) + non-protein part (cofactor).
- Apoenzyme + cofactor ? **holoenzyme**.

### Based on Location

- **Intracellular enzymes** ? metabolic enzymes.
- **Extracellular enzymes** ? digestive enzymes (amylase, lipase).

### Based on Reaction Rate

- **Constitutive enzymes** ? always present.
  - **Inducible enzymes** ? upregulated when substrate appears (e.g.,  $\beta$ -galactosidase).
- 

## Co-enzymes

---

Co-enzymes are **organic, non-protein molecules** required by some enzymes for catalytic activity.

Most are derived from **vitamins**.

---

## 1. Coenzymes Derived from B-Complex Vitamins

- **NAD<sup>+</sup> / NADP<sup>+</sup> (from Niacin)**

- Participate in **oxidation–reduction** reactions.
- Accept **hydride ions (H<sup>-</sup>)**.
- Used by dehydrogenases (e.g., lactate dehydrogenase).

---

- **FAD / FMN (from Riboflavin)**

- Accept **two hydrogens** in redox reactions.
- Cofactor for **succinate dehydrogenase**.

---

- **Coenzyme A (from Pantothenic Acid)**

- Carries **acyl groups**.
- Essential for **fatty acid oxidation**, Krebs cycle, cholesterol synthesis.

---

- **Pyridoxal Phosphate – PLP (from Vitamin B<sub>6</sub>)**

- Coenzyme for **transamination, decarboxylation, deamination**.
- Used by aminotransferases (ALT, AST).

---

- **Biotin (Vitamin B<sub>7</sub>)**

- Cofactor for **carboxylation** reactions.
- Enzymes: **pyruvate carboxylase, acetyl-CoA carboxylase**.

---

- **Tetrahydrofolate – THF (from Folate)**

- Transfers **one-carbon units** (methyl, formyl).
- Essential for **nucleotide synthesis**.

---

- **Cobalamin Coenzymes (Vitamin B<sub>12</sub>)**

- Required for **methionine synthase** and **methylmalonyl-CoA mutase**.
- 

- **Thiamine Pyrophosphate – TPP (from Vitamin B?)**

- Coenzyme in **oxidative decarboxylation** (PDH,  $\alpha$ -ketoglutarate dehydrogenase).
  - Also for **transketolase** in HMP shunt.
- 

## 2. Non-Vitamin Coenzymes

- **Coenzyme Q (Ubiquinone)**

- Electron carrier in the electron transport chain.

- **Heme**

- Cofactor in cytochromes & peroxidases.

- **Lipoic Acid**

- Coenzyme for **pyruvate dehydrogenase** and  **$\alpha$ -ketoglutarate dehydrogenase**.
- 

## Enzyme Prosthetic Groups vs Coenzymes

---

- **Coenzymes** ? loosely bound, dissociable.
  - **Prosthetic groups** ? tightly or covalently attached.
- 

## Clinically Important Points

---

- B-complex vitamin deficiency ? enzyme dysfunction ? metabolic disorders.
- ALT/AST require **PLP**; deficiency ? defective amino-acid metabolism.
- B<sub>1</sub> deficiency ? PDH dysfunction ? lactic acidosis.

## Mode of Action of Enzymes

---

## 1. Lowering Activation Energy

- Enzymes speed up reactions by lowering **activation energy (E<sub>a</sub>)**.
- They do NOT change **ΔG (free energy)** or **equilibrium constant**.

## 2. Formation of Enzyme–Substrate (ES) Complex

- Substrate binds to the enzyme's **active site** → forms ES complex.
- ES complex stabilizes the transition state → faster product formation.

## 3. Models of Enzyme Action

### • Lock and Key Model

– Active site is rigid and fits substrate exactly.

### • Induced Fit Model

– Active site is flexible; binding induces conformational change.

– More accurate for most enzymes.

## 4. Reaction Pathway



- Enzyme remains unchanged after reaction.

---

## Active Center (Active Site)

---

### Definition

- The region of the enzyme where **substrate binding** and **catalysis** occur.

### Characteristics

- Occupies only a **small portion** of the enzyme.
- Formed by **specific amino acids** (Ser, His, Asp, Cys, Lys, Glu).
- 3D orientation determines specificity.

### Types of Active Site Residues

- **Binding residues** → hold the substrate.
- **Catalytic residues** → perform bond-breaking/bond-making.

## Microenvironment

- Hydrophobic pocket
- Correct orientation for catalysis
- Stabilizes transition state

## Substrate Specificity

- Absolute (urease acts only on urea)
- Group-specific (hexokinase phosphorylates many hexoses)
- Stereo-specific (L-amino acid oxidase)

---

## Enzyme Kinetics (Michaelis–Menten)

---

### Basic Equation

$$v = (V_{\max} \times [S]) / (K_m + [S])$$

### Definitions

- **v** = reaction velocity
- **V<sub>max</sub>** = maximum velocity when enzyme is saturated
- **K<sub>m</sub>** = substrate concentration at  $\frac{1}{2}$  **V<sub>max</sub>**

### Assumptions

- ES complex formation is reversible.
- Steady-state concentration of ES.
- Substrate  $\gg$  enzyme concentration.

### Interpretation

- At **low [S]** ? reaction first-order (rate  $\propto$  [S]).
- At **high [S]** ? reaction zero-order (rate independent of [S]).
- V<sub>max</sub> depends on **enzyme concentration**.
- K<sub>m</sub> is independent of enzyme concentration.

---

## Michaelis Constant (K<sub>m</sub>)

---

## Definition

- $K_m$  is the substrate concentration at which the reaction velocity is **half of  $V_{max}$** .

## Significance

- Measures **enzyme affinity** for substrate.
  - **Low  $K_m$**  ? high affinity ? enzyme saturates quickly
  - **High  $K_m$**  ? low affinity
- GIVES a quantitative measure of how strongly an enzyme binds its substrate.

## Clinical Uses

- Hexokinase has **low  $K_m$**  ? high affinity ? active even at low glucose.
- Glucokinase has **higher  $K_m$**  ? active only after meals ? prevents hypoglycemia.
- Useful in diagnosing genetic enzyme defects.

---

## Enzyme Activation

---

### 1. Zymogen Activation

- Enzymes synthesized in **inactive precursor** forms (zymogens).
- Activated by **proteolytic cleavage**.

Examples:

- Pepsinogen ? pepsin
- Trypsinogen ? trypsin

### 2. Allosteric Activation

- Activator binds to **allosteric site** ? increases enzyme activity.

Example:

- ATP activates phosphofructokinase-1 (PFK-1) in glycolysis (at high energy states).

### 3. Covalent Modification

- **Phosphorylation/dephosphorylation** alters enzyme activity.

Examples:

- Glycogen phosphorylase active when phosphorylated.

- Acetyl-CoA carboxylase active when dephosphorylated.

#### 4. Metal Ion Activation

- Some enzymes require metal ions as activators.

Examples:

- $Mg^{2+}$  kinases
- $Zn^{2+}$  carbonic anhydrase
- $Ca^{2+}$  clotting enzymes

#### 5. pH and Temperature Activation

- Each enzyme has optimum pH & temperature.
- Small changes can enhance activity until denaturation occurs.

### Competitive Inhibition

---

#### Definition

- In competitive inhibition, the **inhibitor resembles the substrate** and **competes for the active site** of the enzyme.

#### Key Mechanism

- Inhibitor binds **only to the active site** of free enzyme (E).
- Prevents ES complex formation.

#### Reversibility

- Reversible by **increasing substrate concentration**.

#### Effect on Kinetics

- **$V_{max}$**  ? *unchanged*
- **$K_m$**  ? *increased* (lower affinity, more substrate needed)
- **Lineweaver–Burk Plot:**



- Lines intersect on the **y-axis** (same  $V_{max}$ ).
- Slope increases.

## Examples

- **Malonate** inhibits succinate dehydrogenase.
- **Statins** competitively inhibit HMG-CoA reductase.
- **Methotrexate** inhibits dihydrofolate reductase.

## Clinical Relevance

- Increasing substrate (e.g., high-dose folate) can overcome methotrexate toxicity.

---

## Noncompetitive Inhibition

---

### Definition

- Inhibitor binds to a **site other than the active site** (allosteric site).
- Binding distorts enzyme conformation ? reduces activity.

### Key Mechanism

- Inhibitor can bind to **E or ES** complex.
- Does **not compete** with substrate.

### Reversibility

- Cannot be reversed by increasing substrate concentration.

### Effect on Kinetics

- **$V_{max}$**  ? *decreased*
- **$K_m$**  ? *unchanged* (affinity same, but active enzyme molecules fewer)
- **Lineweaver–Burk Plot:**
  - Lines intersect on the **x-axis** (same  $K_m$ ).
  - Slope increases, y-intercept increases.

## Examples

- **Cyanide** inhibits cytochrome oxidase.
- **Heavy metals** (Hg<sup>2+</sup>, Ag<sup>+</sup>) inhibit SH-containing enzymes.
- **Alanine** noncompetitively inhibits pyruvate kinase.

## Clinical Relevance

- Removal of inhibitor or chelation of metal ions can restore activity (e.g., BAL for arsenic poisoning).

## Allosteric Inhibition

---

### Definition

- Allosteric inhibition occurs when an inhibitor binds to an **allosteric (regulatory) site**, not the active site.
- Binding causes a **conformational change** → decreased enzyme activity.

### Characteristics

- Does **not resemble the substrate**.
- Can act rapidly and reversibly.
- Often occurs in **regulatory enzymes** of metabolic pathways.
- Shows **sigmoidal (S-shaped) kinetics**, not Michaelis–Menten.

### Example

- ATP inhibits **phosphofructokinase-1 (PFK-1)** in glycolysis.
- CTP inhibits **aspartate transcarbamoylase**.

---

## Key (Regulatory) Enzymes

---

### Definition

- Enzymes that catalyze **rate-limiting steps** of metabolic pathways.

### Properties

- Usually **allosteric enzymes**.
- Irreversible, early in the pathway.
- Highly regulated by activators/inhibitors.

### Important Examples

- **PFK-1** – rate-limiting enzyme of glycolysis
  - **Glutamate dehydrogenase** – amino-acid metabolism
  - **HMG-CoA reductase** – cholesterol synthesis
  - **Glycogen phosphorylase** – glycogen breakdown
  - **Carbamoyl phosphate synthetase I** – urea cycle
  - **Acetyl-CoA carboxylase** – fatty acid synthesis
- 

### Feedback Inhibition

---

#### Definition

- End-product of a metabolic pathway **inhibits the first committed step** ? prevents overproduction.

#### Mechanism

- End-product binds to an **allosteric site** of the initial enzyme.
- Reduces enzyme activity by conformational change.

#### Importance

- Maintains metabolic balance.
- Prevents waste of energy and substrates.
- Quick and reversible control mechanism.

#### Examples

- **Isoleucine inhibits threonine dehydratase.**
  - **Cholesterol inhibits HMG-CoA reductase.**
  - **ATP inhibits PFK-1.**
-

## Uncompetitive Inhibition

---

### Definition

- Inhibitor binds **only to the ES complex**, not to free enzyme.
- Prevents formation of product ? ES becomes ESI (inactive).

### Effect on Kinetics

- **V<sub>max</sub>** ? decreased
- **K<sub>m</sub>** ? decreased

(Because inhibitor locks ES complex, making enzyme appear to have higher affinity)

### Reversibility

- Cannot be reversed by increasing substrate concentration.

### Lineweaver–Burk Characteristics

- Lines are **parallel** (same slope).
- Y-intercept increases; x-intercept shifts.

### Example

- Lithium inhibits **inositol monophosphatase** uncompetitively.

---

## Lineweaver–Burk Plot

---

### Purpose

- Double reciprocal plot used to determine **K<sub>m</sub>** and **V<sub>max</sub>** and to differentiate types of inhibition.

### Equation

$$1/v = (K_m/V_{max}) \times (1/[S]) + 1/V_{max}$$

- Straight line where:
  - **Y-intercept** =  $1/V_{max}$
  - **X-intercept** =  $-1/K_m$

– **Slope** =  $K_m/V_{max}$

## Interpretation in Inhibition

### 1. Competitive Inhibition

- $V_{max}$  same
- $K_m$  increases
- Lines intersect at **y-axis**

### 2. Noncompetitive Inhibition

- $V_{max}$  decreases
- $K_m$  unchanged
- Lines intersect at **x-axis**

### 3. Uncompetitive Inhibition

- $V_{max}$  decreases
- $K_m$  decreases
- Lines are **parallel**

## Advantages

- Easy comparison of inhibition patterns.

## Disadvantages

- Distorts error at low substrate concentrations; Eadie–Hofstee plot is more accurate.

## Covalent Modification

---

### Definition

- Regulation of enzyme activity through **reversible covalent addition or removal** of a chemical group.

### Most Common: Phosphorylation / Dephosphorylation

- **Kinases** add phosphate (ATP → ADP).
- **Phosphatases** remove phosphate.

## Effects

- Can **activate or inhibit** depending on the enzyme.

## Examples

- **Glycogen phosphorylase** ? active when phosphorylated.
- **Glycogen synthase** ? inactive when phosphorylated.
- **Acetyl-CoA carboxylase** ? active when dephosphorylated.

## Other Covalent Modifications

- Adenylation
- Methylation
- ADP-ribosylation
- Ubiquitination (? marks proteins for degradation)

---

## Repression

---

### Definition

- Long-term regulation where synthesis of an enzyme is **suppressed at the gene level** when its product is abundant.

### Characteristics

- Slower, affects **amount** of enzyme, not immediate activity.
- Seen in bacteria and human metabolic pathways.

### Example

- High cholesterol represses **HMG-CoA reductase** gene expression.

---

## Induction

---

### Definition

- Increased **gene expression** ? increased enzyme synthesis in response to a metabolite or drug.

## Examples

- High carbohydrate diet induces **glucokinase**.
- Barbiturates induce **cytochrome P450** enzymes.
- Lactose induces  **$\beta$ -galactosidase** in bacteria.

## Importance

- Allows metabolic adaptation to environmental or dietary conditions.

---

## Factors Affecting Enzyme Activity

---

### 1. Temperature

- Activity increases with temperature up to optimum ( $\sim 37^\circ\text{C}$ ).
- High temperature  $\beta$  denaturation.

### 2. pH

- Each enzyme has an **optimum pH**.
- Extreme pH  $\beta$  denatures enzyme.

### 3. Substrate Concentration

- Activity increases until  **$V_{\text{max}}$**  is reached (enzyme saturation).
- Follows the Michaelis–Menten curve.

### 4. Enzyme Concentration

- Rate  $\beta$  enzyme concentration (when substrate is in excess).

### 5. Product Concentration

- Accumulation of product slows reaction (product inhibition).

### 6. Activators

- Metal ions ( $\text{Mg}^{2+}$ ,  $\text{Zn}^{2+}$ ,  $\text{Ca}^{2+}$ ) often essential.
- Example: kinases need  **$\text{Mg}^{2+}$** .

## 7. Inhibitors

- Competitive, noncompetitive, uncompetitive, allosteric inhibitors decrease activity.
- 

### Isoenzymes (Isozymes)

---

#### Definition

- Different molecular forms of the **same enzyme** that catalyze the same reaction but differ in structure, kinetics, and tissue distribution.

#### Clinical Significance

- Useful in diagnosing **tissue damage** because each isoenzyme is tissue-specific.
- 

### Lactate Dehydrogenase (LDH) Isoenzymes

---

LDH has **five isoenzymes** (tetramers of H and M subunits):

1. **LDH-1 (H<sub>4</sub>)** – Heart, RBC
2. **LDH-2 (H<sub>3</sub>M<sub>1</sub>)** – Reticuloendothelial system
3. **LDH-3 (H<sub>2</sub>M<sub>2</sub>)** – Lungs
4. **LDH-4 (H<sub>1</sub>M<sub>3</sub>)** – Kidneys, pancreas
5. **LDH-5 (M<sub>4</sub>)** – Liver, skeletal muscle

#### Clinical Patterns

- **MI (heart attack)** ? LDH-1 ? above LDH-2 (flipped pattern).
  - **Liver disease / muscle injury** ? LDH-5 ?.
  - **Hemolysis** ? LDH-1 ? (released from RBCs).
- 

### Creatine Kinase (CK) Isoenzymes

---



CK exists in **three isoforms**:

1. **CK-BB (CK-1)**

- Brain, smooth muscle
- Increased in CNS injury

2. **CK-MB (CK-2)**

- Heart muscle
- **Most specific marker for myocardial infarction**
- Rises 4–6 hours after MI, peaks at 24 hours, normal in 48 hours

3. **CK-MM (CK-3)**

- Skeletal muscle
- Increased in muscular dystrophy, rhabdomyolysis, trauma

---

## Clinical Use

---

- LDH isoenzymes ? differentiate liver, heart, lung, muscle diseases.
- CK-MB ? early diagnosis of **acute myocardial infarction**.
- CK-BB ? stroke, CNS tumors.
- CK-MM ? muscle injury.

## Specificity of Enzymes

---

Enzymes show **high specificity** toward substrates and reactions.

### 1. Absolute Specificity

- Enzyme acts only on **one substrate**.
- Example: **Urease** ? only urea.

### 2. Group Specificity

- Acts on substrates with **similar functional groups**.
- Example: **Hexokinase** ? phosphorylates many hexoses.

### 3. Bond Specificity

- Acts only on a particular type of **bond**.
- Example: **Esterases** ? hydrolyze ester bonds.

### 4. Stereospecificity

- Distinguish between **D- and L-forms**.
- Example: **L-amino acid oxidase, D-lactate dehydrogenase**.

### 5. Reaction Specificity

- One type of chemical transformation only.
- Example: **Oxidoreductases** ? redox reactions only.

---

## Enzyme Engineering

---

### Definition

Modification of enzymes through biochemical, genetic, or structural changes to improve function.

### Methods

- **Site-directed mutagenesis** ? change specific amino acids.
- **Directed evolution** ? repeated mutation + selection.
- **Fusion proteins** ? catalytic domain + tag (His-tag).

### Applications

- Improved stability (heat-stable enzymes).
  - Reduced inhibition.
  - Faster industrial biocatalysis (detergent enzymes).
  - Design of **insulin analogs**, engineered proteases, and enzyme replacement therapies.
-

## Enzyme Units

---

### 1. International Unit (IU)

- Amount of enzyme that converts **1 micromole of substrate per minute** under defined conditions.

### 2. Katal

- SI unit.
- Amount converting **1 mole of substrate per second**.
- (1 katal = 60,000 IU)

### 3. Specific Activity

- Units of enzyme **per mg of protein**.
- Indicates enzyme **purity**.

### 4. Turnover Number (kcat)

- Number of substrate molecules converted to product **per enzyme molecule per second**.

---

## Isoenzymes (Isozymes)

---

*(Already partly covered earlier, expanded here)*

### Definition

Different molecular forms of the **same enzyme** with:

- same catalytic action,
- different amino acid sequences,
- different tissue distribution.

### Example Families

- **LDH** (LDH-1 ? LDH-5)
- **Creatine Kinase** (CK-BB, CK-MB, CK-MM)
- **Alkaline phosphatase (ALP)** isoenzymes – liver, bone, placenta
- **Amylase** – pancreatic vs salivary

---

## Diagnostic Enzymes (Clinical Enzymology)

---

Enzymes used as **biomarkers** for tissue injury.

### 1. Cardiac Enzymes

- **CK-MB** ? MI (rises 4–6 h, normal in 48 h)
- **LDH-1** ? MI (LDH1 > LDH2 = flipped pattern)
- **Troponin** (not an enzyme but key marker)

### 2. Liver Enzymes

- **ALT (SGPT)** ? hepatocellular damage
- **AST (SGOT)** ? liver & muscle
- **ALP** ? cholestasis, bone disease
- **GGT** ? alcoholism, biliary obstruction

### 3. Pancreatic Enzymes

- **Amylase**
- **Lipase** ? more specific for acute pancreatitis

### 4. Muscle Enzymes

- **CK-MM** ? muscle injury, rhabdomyolysis
- **Aldolase** ? muscle diseases

### 5. Bone/Placenta

- **Bone ALP** ? rickets, Paget disease
- **Placental ALP** ? pregnancy, germ cell tumors

---

## Isoenzyme Electrophoresis

---

### Definition

Separation of isoenzymes based on differences in **charge, mobility, and size**.

### Methods

- Agarose gel electrophoresis
- Cellulose acetate electrophoresis
- Isoelectric focusing

### LDH Example

LDH isoforms migrate differently:

- LDH-1 (H4) ? fastest, most negative, moves furthest
- LDH-5 (M4) ? slowest, least negative

### Clinical Uses

- Diagnosing **myocardial infarction** (LDH1 > LDH2).
- Differentiating **liver vs bone ALP**.
- Identifying cancer-related isoenzyme patterns.
- Confirming **salivary vs pancreatic** amylase.

---

### Summary (Exam-Ready One-Liners)

---

- Enzymes show **absolute, group, stereo, bond** specificity.
- Enzyme engineering modifies catalytic efficiency and stability.
- IU = amount of enzyme converting **1 ?mol/min**.
- Isoenzymes differ in structure but catalyze **same reaction**.
- LDH-1 elevation ? myocardial infarction.
- CK-MB ? most specific enzymatic marker for MI.
- ALP high with GGT normal ? **bone disease**.
- Electrophoresis separates isoenzymes based on charge differences.

---

### Frequently Asked Questions (FAQs)

---

#### 1. What determines enzyme specificity?

The **3D structure of the active site**, which recognizes the substrate based on shape, charge, and stereochemistry.

---

## 2. What is absolute specificity?

Enzyme acts on **only one substrate**. Example: **Urease ? Urea**.

---

## 3. What is group specificity?

Enzyme acts on substrates with **similar functional groups** (e.g., hexokinase).

---

## 4. What is the purpose of enzyme engineering?

To improve enzyme **stability, activity, or specificity** using genetic or chemical modifications.

---

## 5. What is site-directed mutagenesis?

A technique to modify **specific amino acids** in an enzyme to alter function.

---

## 6. What is an international unit (IU) of enzyme?

Amount of enzyme that converts **1 micromole of substrate per minute**.

---

## 7. What is a Katal?

SI unit of enzyme activity = **1 mole of product per second**.

---

## 8. What is specific activity?

Units of enzyme per **mg of protein**—an indicator of enzyme **purity**.

---

## 9. What is an isoenzyme?

Different molecular forms of an enzyme, with same function but **different structure and tissue distribution**.

---

## 10. Why are isoenzymes clinically important?

They help identify **which tissue is damaged** during disease.

---

**11. Which LDH isoenzyme indicates myocardial infarction?**

**LDH-1 > LDH-2** (“flipped pattern”).

---

**12. Which CK isoenzyme is specific for heart muscle?**

**CK-MB.**

---

**13. When does CK-MB rise after an MI?**

Rises at **4–6 hours**, peaks at 24 hours, normal in 48 hours.

---

**14. Which enzymes rise in acute pancreatitis?**

**Amylase and lipase**, with lipase being more specific.

---

**15. What enzyme pattern suggests liver cell damage?**

? **ALT**, ? **AST** (AST may rise higher in alcohol-related damage).

---

**16. What enzyme pattern suggests biliary obstruction?**

? **ALP** and ? **GGT**.

---

**17. What does increased bone ALP indicate?**

Rickets, osteomalacia, Paget disease.

---

**18. Which isoenzyme increases in skeletal muscle injury?**

**CK-MM.**

---

**19. How are isoenzymes separated?**

By **electrophoresis** (agarose gel, cellulose acetate) or **isoelectric focusing**.

---

20. Which amylase isoenzyme rises in acute pancreatitis?

Pancreatic amylase.

---

21. What is repression in enzyme regulation?

Reduction in **enzyme synthesis at gene level** when product is abundant.

---

22. What is induction?

Increased **enzyme synthesis** in response to a substrate, hormone, or drug.

---

23. Give an example of an induced enzyme.

Cytochrome P450 enzymes induced by **barbiturates**.

---

24. Which metal ion activates most kinases?

**Mg<sup>2+</sup>**.

---

25. What is the advantage of isoenzyme electrophoresis?

It distinguishes tissue-specific enzyme forms, aiding **diagnosis** (heart vs liver vs bone pathology).

## MCQs

---

1. Absolute specificity is seen in which enzyme?

- A. Hexokinase
- B. Trypsin
- C. Urease
- D. Lipase

**Answer: C**



---

**2. Hexokinase shows which type of specificity?**

- A. Absolute
- B. Group
- C. Bond
- D. Reaction

**Answer: B**

---

**3. Stereospecificity is shown by:**

- A. Pepsin
- B. D-amino acid oxidase
- C. Amylase
- D. Catalase

**Answer: B**

---

**4. Site-directed mutagenesis is used in:**

- A. Enzyme repression
- B. Enzyme engineering
- C. Feedback inhibition
- D. Zymogen activation

**Answer: B**

---

**5. International Unit of enzyme activity means:**

- A. 1 mmol/min
- B. 1  $\mu$ mol/min
- C. 1 mol/sec
- D. 1  $\mu$ mol/sec

**Answer: B**

---

**6. Specific activity indicates:**

- A. Purity of enzyme
- B. pH of enzyme
- C. Amount of substrate
- D. Temperature stability

**Answer: A**

---

**7. Which isoenzyme is elevated in myocardial infarction?**

- A. LDH-5
- B. LDH-3
- C. LDH-1
- D. LDH-4

**Answer: C**

---

**8. LDH-1 > LDH-2 pattern is called:**

- A. Forward pattern
- B. Flipped pattern
- C. Reverse pattern
- D. Saturation pattern

**Answer: B**

---

**9. CK-MB is a marker for:**

- A. Liver failure
- B. Acute pancreatitis
- C. Skeletal muscle injury
- D. Myocardial infarction

**Answer: D**

---

**10. Which enzyme rises earliest after MI?**

- A. LDH
- B. CK-MB
- C. Troponin I

D. AST

**Answer: B**

---

**11. Which enzyme rises highest in obstructive jaundice?**

A. ALT

B. AST

C. ALP

D. LDH

**Answer: C**

---

**12. GGT elevation indicates:**

A. Acute bone disease

B. Alcoholic liver disease

C. Muscle injury

D. Rickets

**Answer: B**

---

**13. Which enzyme is most specific for acute pancreatitis?**

A. Amylase

B. Trypsin

C. Lipase

D. Elastase

**Answer: C**

---

**14. Bone ALP is elevated in:**

A. Cirrhosis

B. Paget disease

C. Myocardial infarction

D. Cushing syndrome

**Answer: B**

---

**15. CK-BB is mainly found in:**

- A. Heart
- B. Skeletal muscle
- C. Brain
- D. Liver

**Answer: C**

---

**16. Isoenzymes differ in:**

- A. Function
- B. Activation energy
- C. Amino acid sequence
- D. Reaction catalyzed

**Answer: C**

*(Reaction catalyzed is same.)*

---

**17. Isoenzymes are best separated by:**

- A. Simple centrifugation
- B. Electrophoresis
- C. Precipitation
- D. Dialysis

**Answer: B**

---

**18. In non-competitive inhibition, which parameter changes?**

- A.  $K_m$  increases
- B.  $K_m$  decreases
- C.  $V_{max}$  decreases
- D.  $V_{max}$  increases

**Answer: C**

---

**19. An enzyme showing sigmoidal kinetics is usually:**

- A. A simple enzyme
- B. An allosteric enzyme
- C. A hydrolase
- D. A zymogen

**Answer: B**

---

**20. Feedback inhibition usually acts on:**

- A. The last enzyme of the pathway
- B. Any random enzyme
- C. The rate-limiting enzyme
- D. The fastest enzyme

**Answer: C**

---

**21. An inhibitor that binds only to the ES complex is:**

- A. Competitive
- B. Non-competitive
- C. Uncompetitive
- D. Allosteric

**Answer: C**

---

**22. Which enzyme requires  $Mg^{2+}$  for activation?**

- A. Pepsin
- B. Kinases
- C. Urease
- D. Lipase

**Answer: B**

---

**23. Enzyme induction means:**

- A. Increase in substrate concentration
- B. Increase in enzyme activity
- C. Increase in enzyme synthesis

D. Decrease in enzyme affinity

**Answer: C**

---

**24. Cytochrome P450 enzymes are induced by:**

- A. Vitamin C
- B. Barbiturates
- C. Insulin
- D. Iron deficiency

**Answer: B**

---

**25. Enzyme repression occurs when:**

- A. Substrate is in excess
- B. Product accumulates
- C. Temperature increases
- D. pH increases

**Answer: B**

## Viva Voce

---

**1. What is enzyme specificity?**

It is the ability of an enzyme to choose a **single substrate or group of substrates** based on its active-site structure.

---

**2. What is absolute specificity?**

The enzyme acts on **only one specific substrate**.

Example: Urease ? urea.

---

**3. What is group specificity?**

The enzyme catalyzes reactions of substrates with **similar functional groups**.

Example: Hexokinase.

---

#### 4. What is stereospecificity?

The enzyme distinguishes between **D- and L-forms** of molecules.

---

#### 5. What is reaction specificity?

The enzyme catalyzes only **one type of chemical reaction**, regardless of substrate variety.

---

#### 6. What is enzyme engineering?

Modification of enzyme structure by **genetic or chemical methods** to improve activity, stability, or specificity.

---

#### 7. Give an example of enzyme engineering.

Site-directed mutagenesis to create **heat-stable enzymes**.

---

#### 8. What is site-directed mutagenesis?

Technique to change a **specific amino acid** in a protein to alter its function.

---

#### 9. What is an International Unit (IU)?

Amount of enzyme that converts **1 μmol of substrate per minute**.

---

#### 10. What is a Katal?

SI unit of enzyme activity (1 mole of substrate converted per second).

---

#### 11. What is specific activity?

Enzyme units **per mg of protein**—indicator of enzyme purity.

---

**12. What is turnover number (kcat)?**

The number of **substrate molecules converted per enzyme molecule per second**.

---

**13. What are isoenzymes?**

Different molecular forms of the **same enzyme** with different structures but identical catalytic function.

---

**14. Why are isoenzymes important clinically?**

They help identify **which tissue is damaged**, since each isoenzyme is tissue-specific.

---

**15. Which LDH isoenzyme rises in myocardial infarction?**

**LDH-1**, showing the *flipped pattern* (LDH-1 > LDH-2).

---

**16. Which CK isoenzyme is specific for cardiac muscle?**

**CK-MB**.

---

**17. When does CK-MB appear after MI?**

Rises in 4–6 hours, peaks at 24 hours, normal after 48 hours.

---

**18. Which enzyme is most specific for acute pancreatitis?**

**Lipase**.

---

**19. Which enzyme rises in cholestasis?**

**ALP**, along with **GGT**.

---

**20. Which ALP isoenzyme rises in bone disease?**

**Bone ALP**.

---



**21. What does elevated CK-MM indicate?**

Skeletal muscle injury or rhabdomyolysis.

---

**22. What is enzyme electrophoresis?**

A technique to separate isoenzymes based on **charge and mobility**.

---

**23. Which technique gives best isoenzyme separation?**

**Isoelectric focusing**, based on pI differences.

---

**24. What is feedback inhibition?**

The end-product inhibits the **rate-limiting enzyme** of its own pathway.

---

**25. What is repression?**

Decreased **enzyme synthesis** at the gene level due to excess product.

---

**26. What is induction?**

Increased **enzyme synthesis** in response to a metabolite, hormone, or drug.

---

**27. What type of regulation do allosteric enzymes show?**

**Sigmoidal kinetics** and rapid, reversible control.

---

**28. What is the major allosteric inhibitor of PFK-1?**

**ATP**.

---

**29. Which metal ion activates most kinases?**

**Mg<sup>2+</sup>**.

---

**30. What does high GGT with high ALP indicate?**

Obstructive or alcoholic liver disease.